Neutrinos in Large-scale structure

Marilena LoVerde
University of Chicago

Neutrinos in Large-scale structure

Marilena LoVerde
University of Chicago

ML and Zaldarriaga 1310.6459

ML 1405.4855

ML 1404:4858

ML in prep.

Outline

- Neutrinos
- Neutrinos in Cosmology
- Massive Neutrinos in Large-scale structure
 - Halo bias
 - Halo abundance
 - Neutrino halos
- Observational Consequences

Neutrinos


Neutrinos


flavor eigenstates

Velectror

Vtau

Vmuon






flavor eigenstates z mass eigenstates

Neutrinos Plavor eigenstates






oscillation data gives mass splittings

$$m_2^2 - m_1^2 = (7.5 \pm 0.2) 10^{-5} \text{ eV}^2$$

(solar neutrino oscillations)

$$|m_3^2 - m_2^2| = (2.32^{+0.12}_{-0.08}) 10^{-3} \text{ eV}^2$$

(atmospheric neutrino oscillations)







but the absolute masses are unknown!

``Normal"

mass

_____v₃ ≈ 0.049eV _____v₂ ≈ 0.009eV v₁ ≈ 0eV






but the absolute masses are unknown!

mass






but the absolute masses are unknown!



Absolute scale, hierarchy

Tritium β decay: $m_{\nu_{\overline{e}}} \lesssim 2eV$ upper bound on mass (Troitsk experiment 2011)



Big Questions

- Where do neutrino masses come from?
- What is the absolute mass scale?
- Are there additional sterile neutrinos?
- Can neutrinos generate matter-antimatter asymmetry of the universe?

Big Questions

- Where do neutrino masses come from?
- · What is the absolute mass scale?
- · Are there additional sterile neutrinos?
- Can neutrinos generate matter-antimatter asymmetry of the universe?

Big Questions

- Where do neutrino masses come from?
- · What is the absolute mass scale?
- Are there additional sterile neutrinos?
- Can neutrinos generate matter-antimatter asymmetry of the universe?

Astrophysics has previously informed neutrino physics! (solar neutrino problem \Longrightarrow neutrino oscillations)

Neutrinos in Cosmology

e+ e- annihilation



hotter


big bang nucleosynthesis

galaxy formation

today! 14 billion years time
10° years colder

T ~ 0.2 eV photons decoupled

e+ e- annihilation




hotter


T ~ 1 MeV big bang nucleosynthesis

neutrinos in equilibrium with photons e +, e-

$$T_{\gamma} = T_{\nu} \propto 1/a$$



galaxy formation



109 years



today! 14 billion years time colder

T ~ 0.2 eV photons

decoupled

e+ e- annihilation




hotter


T ~ 1 MeV big bang nucleosynthesis

neutrinos in equilibrium with photons e


neutrinos decoupled Tv ∝ 1/a

$$T_{\gamma} = T_{\nu} \propto 1/a$$





galaxy formation



109 years



today! 14 billion years time colder

T ~ 0.2 eV photons decoupled

e+ e- annihilation

galaxy formation

109 years



today! 14 billion years time colder

hotter

T ~ 1 MeV big bang nucleosynthesis

neutrinos in equilibrium with photons e

+, e- $T_{\gamma} = T_{\nu} \propto 1/a$

neutrinos decoupled

Tv ∝ 1/a

decouple while relativistic

$$n_{1\nu} \sim T_{\nu}^3$$

$$\rho_{1\nu} \sim T_{\nu}^4$$

T ~ 0.2 eV photons decoupled

e+e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$

galaxy formation



109 years

today! 14 billion years time

colder

hotter

T ~ 1 MeV big bang nucleosynthesis

neutrinos in equilibrium

decoupled

Tv ∝ 1/a

with photons e +, e-

 $T_{\gamma} = T_{\nu} \propto 1/a$

neutrinos

decouple while relativistic

$$n_{1\nu} \sim T_{\nu}^3$$

$$\rho_{1\nu} \sim T_{\nu}^4$$

T ~ 0.2 eV photons

decoupled

$$T_{\gamma} \approx \frac{11}{4} T_{\nu} \qquad n_{1\nu} \approx 100/\text{cm}^{3}$$

$$T_{\nu} \approx 0.17 \text{ meV}$$

e+ e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$





hotter

T ~ 1 MeV big bang nucleosynthesis

neutrinos in equilibrium with photons e +, e-


$$T_{\gamma} = T_{\nu} \propto 1/a$$

$$n_{1\nu} \sim T_{\nu}^3$$

$$\rho_{1\nu} \sim T_{\nu}^{4}$$

$$T_{\gamma} \approx \left(\frac{11}{4}\right)^{1/3} T_{\nu}$$

galaxy formation






109 years



today! 14 billion years time colder

T ~ 0.2 eV photons decoupled




e+ e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$

shi vi

hotter

T ~ 1 MeV big bang nucleosynthesis

galaxy formation



109 years



today! 14 billion years time colder

T ~ 0.2 eV photons decoupled

Once $T_{\nu} << m_{\nu}$ energy density is just $\rho_{\nu} \approx \Sigma m_{\nu} n_{\nu}$

e+ e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$

galaxy formation



hotter

T ~ 1 MeV big bang nucleosynthesis

109 years


today! 14 billion years time colder

T ~ 0.2 eV photons decoupled

Once $T_{\nu} << m_{\nu}$ energy density is just $\rho_{\nu} \approx \Sigma m_{\nu} n_{\nu}$

 $n_{1\nu}$ is known, so a measurement of ρ_{ν} gives Σ m_{ν}

e+ e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$





colder

hotter ____

T ~ 1 MeV big bang nucleosynthesis

T ~ 0.2 eV photons

decoupled

Once $T_{\nu} << m_{\nu}$ energy density is just $\rho_{\nu} \approx \sum_{i} m_{\nu} n_{\nu}$

109 years

 $n_{1\nu}$ is known, so a measurement of ρ_{ν} gives Σ m_{ν}

Cosmology at later times, once Tu << mu, tests \(\Sigma \) mu

e+ e- annihilation, T_{γ} boosted relative to T_{ν} , $T_{\gamma} \propto 1/a$

galaxy formation



hotter

T ~ 1 MeV big bang nucleosynthesis

10° years

today! 14 billion years time colder

T ~ 0.2 eV photons decoupled

Once $T_{\nu} << m_{\nu}$ energy density is just $\rho_{\nu} \approx \Sigma m_{\nu} n_{\nu}$


 $n_{1\nu}$ is known, so a measurement of ρ_{ν} gives Σ m_{ν}


Cosmology at later times, once Tu << mu, tests \(\Sigma \) mu




Large-scale Structure

Neutrinos in Large-scale structure







time


(Kravtsov)

The gravitational evolution of large-scale structure is different for fast and slow moving particles

The gravitational evolution of large-scale structure is different for fast and slow moving particles



(clump easily)




(don't clump easily)

The gravitational evolution of large-scale structure is different for fast and slow moving particles



(clump easily)

baryons and cold dark matter



(don't clump easily)

neutrinos (or other exotic light dark matter)

small-scale density
perturbations don't retain
neutrinos

cold dark matter and baryons density perturbation growing


neutrino density perturbation decaying

<u>δρν</u> ρν

small-scale density
perturbations don't retain
neutrinos


cold dark
matter,
baryons and
neutrinos
growing
together

large-scale density perturbations do retain neutrinos










small-scale density perturbations don't retain neutrinos

Growth of matter perturbations is scale-dependent

large-scale density perturbations do retain neutrinos








small-scale density perturbations don't retain neutrinos

large-scale density perturbations do retain neutrinos


Growth of matter perturbations is scale-dependent



Relevant scale:

Typical distance a neutrino can travel in a Hubble time $\lambda_{fs} \sim u_{\nu}/H$





time

small-scale density perturbations don't retain neutrinos

large-scale density perturbations do retain neutrinos

Growth of matter perturbations is scale-dependent

Relevant scale:


Typical distance a neutrino can travel in a Hubble time $\lambda_{fs} \sim u_{\nu}/H$




"free-streaming scale"

time


Scale-dependent growth






Scale-dependent growth





cosmological constraints!



Bond, Efstathiou, Silk 1980 Hu, Eisenstein, Tegmark 1998

Some observable consequences:

Suppressed matter power spectrum (test via lensing, galaxy power spectra)

- Suppressed matter power spectrum (test via lensing, galaxy power spectra)
- Fewer massive halos hosting galaxy clusters

- Suppressed matter power spectrum (test via lensing, galaxy power spectra)
- Fewer massive halos hosting galaxy clusters
 - (But neutrino halos eventually accumulate around the cold dark matter halos!)

- Suppressed matter power spectrum (test via lensing, galaxy power spectra)
- Fewer massive halos hosting galaxy clusters
 - (But neutrino halos eventually accumulate around the cold dark matter halos!)
 - Scale-dependent halo bias

Some observable consequences:

- Suppressed matter power spectrum (test via lensing, galaxy power spectra)
- Fewer massive halos hosting galaxy clusters
 - (But neutrino halos eventually accumulate around the cold dark matter halos!)

(ML & Zaldarriaga 2013)

Scale-dependent halo bias

Some observable consequences:


- Suppressed matter power spectrum (test via lensing, galaxy power spectra)
- Fewer massive halos hosting galaxy clusters
 - (But neutrino halos eventually accumulate around the cold dark matter halos!)


(ML & Zaldarriaga 2013)

Scale-dependent halo bias




Scale-dependent halo bias from massive neutrinos







(these nodes are cold dark matter halos)



(Kravtsov)




(these nodes are cold dark matter halos)




i.e. gravitationally bound blobs of dark matter where galaxies and clusters of galaxies live

(Kravtsov)



of halos is modulated by long-wavelength fluctuations in the matter density field




of halos is modulated by long-wavelength fluctuations in the matter density field


$$\frac{\delta n}{n} \equiv b \frac{\delta \rho}{\rho}$$

b is the halo bias

In a universe with CDM only, the linear evolution of matter fluctuations is independent of their wavelength




In a universe with CDM only, the linear evolution of matter fluctuations is independent of their wavelength



halos can't tell the wavelength of the background matter density perturbation

In a universe with CDM only, the linear evolution of matter fluctuations is independent of their wavelength




halos can't tell the wavelength of the background matter density perturbation



the effect of $\frac{\delta p}{\rho}$ on the halo field (the linear bias) is independent of k

In a universe with CDM only, the linear evolution of matter fluctuations is independent of their wavelength




halos can't tell the wavelength of the background matter density perturbation




the effect of $\frac{\delta p}{\rho}$ on the halo field (the linear bias) is independent of k massive neutrinos break this

halo bias can depend on k




halos neutrinos cold dark matter

WANT: estimate of k-dependence of the halo bias caused by massive neutrinos



halos neutrinos cold dark matter

WANT: estimate of k-dependence of the halo bias caused by massive neutrinos



halos neutrinos cold dark matter


The scale-dependent growth of density perturbations causes halo bias to be scale dependent

The scale-dependent growth of density perturbations causes halo bias to be scale dependent

$$\frac{\delta n}{n}(k) = b \frac{\delta \rho_m}{\rho_m}(k) \qquad \frac{\delta n}{n}(k) = b(k) \frac{\delta \rho_m}{\rho_m}(k)$$


Prescription for calculating the halo bias in a universe with massive neutrinos


initial density field




initial density field









initial density field



initial proto-halo distribution





late time halo distribution




initial density field

initial proto-halo distribution











initial density field

initial proto-halo distribution






 $\frac{\delta n}{n} \equiv \boxed{b} \frac{\delta \rho}{\rho}$ | long-wavelength


late time halo distribution


want this!



initial density field

initial proto-halo distribution





(i) In this step, the scale-dependent evolution of density perturbations causes

$$\frac{\delta n_{\text{Lagrangian}}(k) = b_{\text{Lagrangian}}(k) \frac{\delta \rho_{\text{cdm}}}{\rho_{\text{cdm}}}(k)$$


initial proto-halo distribution

(ii) In this step, the free streaming of neutrinos causes

$$b(k) = \frac{\langle \delta_n(k)\delta_m(k)\rangle}{\langle \delta_m(k)\delta_m(k)\rangle}$$

=
$$(1 + b_{Lagrangian}(k)) \langle \frac{\delta_{cdm}(k)\delta_{m}(k)}{\delta_{m}(k)\delta_{m}(k)} \rangle$$

late time halo distribution



initial proto-halo distribution


(ii) In this step, the free streaming of neutrinos causes

$$b(k) = \frac{\langle \delta_n(k)\delta_m(k)\rangle}{\langle \delta_m(k)\delta_m(k)\rangle}$$

=
$$(1 + b_{Lagrangian}(k)) \langle \frac{\delta_{cdm}(k)\delta_{m}(k)}{\delta_{m}(k)\delta_{m}(k)} \rangle$$

late time halo distribution

(i.e. halos trace CDM —> bias w.r.t total matter is scale-dependent)



initial proto-halo distribution


(ii) In this step, the free streaming of neutrinos causes

$$b(k) = \frac{\langle \delta_n(k) \delta_m(k) \rangle}{\langle \delta_m(k) \delta_m(k) \rangle}$$

=
$$(1 + b_{Lagrangian}(k)) \langle \frac{\delta_{cdm}(k)\delta_{m}(k)}{\delta_{m}(k)\delta_{m}(k)} \rangle$$

late time halo distribution


(i.e. halos trace CDM —> bias w.r.t total matter is scale-dependent)




see also Villaescusa-Navarro, Marulli, Viel, Branchini, Castorina 2013 Castorina, Sefusatti, Sheth, Villaescusa-Navarrow, Viel 2014 Biagetti, Desjacques, Kehagias, Riotto 2014

Numerical estimates for scaledependent halo bias


Scale-dependent changes to δ_{crit}





scale-dependent change to


$$\delta n(k) = \frac{\partial n}{\partial \delta_{crit}} \frac{\partial \delta_{crit}}{\partial \delta_{cdm}(k)} \delta_{cdm}(k)$$



(Use Bhattacharya et al 2011 for n(M| δ_{crit})

scale-dependent change to

$$\delta n(k) = \frac{\partial n}{\partial \delta_{crit}} \frac{\partial \delta_{crit}}{\partial \delta_{cdm}(k)} \delta_{cdm}(k)$$



(Use Bhattacharya et al 2011 for n(M| δ_{crit})

(ML 2014)

scale-dependent change to final bias

 $\delta n(k)/n = b(k) \delta_{matter}(k)$

 $b(k) = \sqrt{P_{hh}(k)/P_{mm}(k)}$


Eulerian halo bias fractional change in

M=1e+13, z=0.501.08 1 massive $\nu m_{\nu} = 0.05 eV$ 1 massive $\nu m_{\nu} = 0.1 eV$ 1 massive $\nu \, m_{\nu} \! = \! 0.2 eV$ 1 massive $\nu m_{\nu} = 0.3 eV$ 3 massive $\nu m_{\nu} = 0.1 eV$ massless 1.04 1.02 1.00 0.98^L 10⁻⁴ 10⁻³ 10⁻² 10^{-1} wavenumber k (Mpc⁻¹)

(Use Bhattacharya et al 2011 for n(M| δ_{crit})

scale-dependent change to final bias


$$\delta n(k)/n = b(k) \delta_{matter}(k)$$




(Use Bhattacharya et al 2011 for n(M| δ_{crit})

scale-dependent change to final bias

$$\delta n(k)/n = b(k) \delta_{matter}(k)$$






Scale-dependent bias

Observational consequences of scale-dependent bias?

Observational consequences of scale dependent bias?


matter power spectrum, or (incorrectly) assuming constant bias

suppression in galaxy power spectrum <u>less</u> than in matter power spectrum




Scale-dependent bias

But the scale-dependent halo bias is itself an observable!




The scale-dependent halo bias is an observable!



$$\sigma_{b1/b2} \sim \frac{1}{\sqrt{n_1 P_{g2g2}}}$$

The scale-dependent halo bias is an observable!





$$\sigma_{b1/b2} \sim \frac{1}{\sqrt{N_{\ell} n_1 C_{g2g2}}}$$

Fewer massive halos from massive neutrinos

initial density field


mass that collapses into halos







initial density field

mass that collapses into halos












Fewer large-amplitude density fluctuations




initial density field













And, higher threshold density to form a halo






using Bhattacharya et al fitting formula for n(M) with our δ_{crit}







Neutrinos accrete onto these cold dark matter halos

escape velocity of dark matter halos




low velocity neutrinos end up bound in halos






low velocity neutrinos end up bound in halos


shown are $m_{\nu} = 0.05 eV$ neutrinos around $M = 10^{14}$ M_{sun} halo






the accreted neutrino
halo is puffy in
comparison with the
CDM!


neutrino mass around halo (Msun)



fraction that's bound to the halo



Neutrino Effects in Large-scale Structure



Accuracy of these predictions?

N-body simulations are the community standard for cold dark matter structure.


Simulations with massive neutrinos?

- (i) Tricky. very few exist, very new
- (ii) Want a model that provides insight into the physical processes responsible for new effects (iii) Don't want to rerun for every possible neutrino
- mass hierarchy scenario
- (iv) It will be great to make comparisons in the future!


Scale-dependent bias from massive neutrinos

comparison with sims looks reasonable!

my calculations







Conclusions

- © Cosmology provides interesting information about neutrinos!
- Scale-dependent halo bias is a new signal of massive neutrinos in large-scale structure
- Scale-dependent halo bias is a new systematic for massive neutrinos in large-scale structure
- Neutrinos decrease abundance of halos Need to be careful with predictions for m_ν ≥ 0.3eV!
- Eventually neutrinos accrete onto very massive halos!